A61K39/00



## WORLD INTELLECTUAL PROPERTY ORGANIZATION International Bureau



#### INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

	INTERNATIONAL APPLICATION PUBLIS	HED	UNDER THE PATENT COOPERATION TREATY (PCT)
	(51) International Patent Classification 4:		(11) International Publication Number: WO 89/12455
	A61K 37/02, 39/04	A1	(43) International Publication Date: 28 December 1989 (28.12.89)
. \ <del>{</del>	(21) International Application Number: PCT/US (22) International Filing Date: 15 June 1989		patent), CH (European patent), DE (European patent), FR (European patent), GB (European patent), IT (European patent), JP, LU (European patent), NL (European
,	(30) Priority data: 207,298  15 June 1988 (15.06.88)  (71) Applicants: WHITEHEAD INSTITUTE FOR BI CAL RESEARCH [US/US]; Nine Cambridg Cambridge, MA 02142 (US). MEDICAL RES COUNCIL [GB/GB]; 20 Mount Pleasant, Lone 4AL (GB).  (72) Inventors: YOUNG, Richard, A.; 5 Sawmill Bro Winchester, MA 01890 (US). YOUNG, Dou Lawnclose Ruislip, Middlesex HA4 6ED (GB)	OMEI e Cent SEARC ion W	claims and to be republished in the event of the receipt of amendments.  N  N  N  N  N  N  N  N  N  N  N  N  N
	(74) Agents: GRANAHAN, Patricia et al.; Hamilton Smith & Reynolds, Two Militia Drive, Lexing 02173 (US).		
DC	C		

(54) Title: STRESS PROTEINS AND USES THEREFOR

#### (57) Abstract

Stress proteins and their use to immunize an individual against a nonviral infection or to induce immune tolerance in an individual, as well as a method of immunizing an individual by administering a selected stress protein and a method of inducing immune tolerance in an individual by administering a selected stress protein.

CALLEY OF STATE OF STATE OF THE STATES

Could lead to the the the lease party to the FO leader to the grouphlies party and the company of the first meaning and and and a contract the first

AN3

## FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

			•	•	
AT	Austria	FI	Finland	ML	Mali
AU	Australia	FR	France	MR	Mauritania
BB	Barbados	GA	Gabon	MW	Malawi
BE	Belgium	G8	United Kingdom	NL	Netherlands
BF	Burkina Fasso	HU	Hungary	NO	Norway
BG	Bulgaria	n	italy	RO	Romania
BJ	Benin	JP	Japan	50	Sudan
BR	Brazil	KP	Democratic People's Republic	SE	Sweden
CF	Central A frican Republic		of Korea	SN	Senegal
CG	Congo	KR	Republic of Korea	SU	Soviet Union
CH	Switzerland	u	Liechtenstein	TD	Chad -
CM	Cameroon	LK	Sri Lanka	TG	Togo
DE A	Germany, Federal Republic of	w	Luxembourg	us	United States of America
OK ES	Denmark	MC	Monaco		Office States of Afficing
<b>F E</b> **	Spain	MG	Madagascar		

#### STRESS PROTEINS AND USES THEREFOR

#### Description

#### Background of the Invention

Although the function of stress proteins is not 05 entirely clear, it appears that some participate in assembly and structural stabilization of certain cellular and viral proteins, and their presence at high concentrations may have an additional stabilizing effect during exposure to adverse conditions. Neidhardt, F.C. 10 and R.A. VanBogelen, In: Escherichia coli and Salmonella typhimurium, Cellular and Molecular Biology, (eds. Neidhardt, F.C., Ingraham, J.L., Low, K.B., Magasanik, B. Schaechter, M. and Umbarger, H.E. (Am. Soc. Microbiol., Washington, D.C.), pp. 1334-1345 (1987); Pelham, H.R.B. Cell, 46:959-961 (1986); Takano, T. and T. Kakefuda, 15 <u>Nature</u>, 239:34-37 (1972); Georgopoulos, C. et al., New Biology, 239:38-41 (1972). Phagocytic host cells produce a hostile environment for foreign organisms, and the ability to produce stress proteins has been implicated in the survival of bacterial pathogens within macrophages 20

Mycobacterium (M.) tuberculosis and Mycobacterium

(M.) leprae are the etiologic agents of tuberculosis and leprosy, respectively. These diseases afflict 20-30

25 million people and continue to present a significant global health problem. Joint International Union Against Tuberculosis and World Health Organization Study Group,

Tubercle, 63:157-169 (1982); Bloom, B. and T. Godal, Rev.

Infect Dis. 5:765-780 (1983). To develop more effective

Christman, M.F. et al., Cell, 41:753-762 (1985).

tools for the diagnosis and prevention of these diseases, it is important to understand the immune response to infection by mycobacterial pathogens.

The antibody and T-cell responses to infection or inoculation with killed mycobacteria have been studied in humans and in animals. Human patients with tuberculosis or leprosy produce serum antibodies directed against at least 12 mycobacterial proteins. Some of these proteins are also recognized by well-characterized murine

- monoclonal antibodies. Mice immunized with mycobacterial lysates produce antibodies that are directed predominantly to six M. tuberculosis and six M. leprae protein antigens. Engers, H.D., Infect. Immun., 48:603-605 (1985); Engers, H.D., Infect. Immun.,
- 51:718-720 (1986). Genes encoding these 12 mycobacterial antigens have been cloned, and recombinant proteins produced from these clones have been used to investigate the human T-lymphocyte response to mycobacterial infection. Husson, R.N. and R.A. Young, <a href="Proc. Natl.">Proc. Natl.</a>
- 20 <u>Acad. Sci., USA</u>, <u>84</u>:1679-1683 (1987); Young, R.A. <u>et al.</u>, <u>Nature</u>, <u>316</u>: 450-452 (1985); Britton, W.J. <u>et al.</u>, <u>Lepr.</u> <u>Rev.</u>, <u>57</u>, Suppl. 2, 67-75 (1986).

Protection against mycobacterial disease involves cell-mediated immunity. Joint International Union

- Against Tuberculosis and World Health Organization Study Group, Tubercle, 63:157-169 (1982); Hahn, H. and S.H.E. Kaufman, Rev. Infect. Dis., 3:1221-1250 (1981). T lymphocytes cloned from patients or from volunteers immunized with killed mycobacteria have been tested for their ability to recognize the recombinant mycobacterial
- proteins. Lymphocyte-proliferation assays demonstrate

that most of the antigens identified with monoclonal antibodies are involved in the T-cell response to mycobacterial infection or vaccination in mice and in humans. Limiting dilution analysis indicates that 20% of the mycobacterial-reactive CD4<sup>+</sup> T lymphocytes in mice immunized with M. tuberculosis recognize a single protein, the 65-kDa antigen. Kaufman, S.H.E. et al., Eur J. Immunol., 17:351-357 (1987).

#### Summary of the Invention

10 The present invention relates to stress proteins and methods of modulating an individual's immune response. either to a pathogen or to his or her own cells, such as occurs in autoimmune diseases. In particular, it relates to the use of such stress proteins as a "vaccine" in immune prophylaxis therapy, which results in an induction or enhancement of immune response to a selected pathogen and as an immunotherapeutic agent in treatment of autoimmune diseases, which results in a decrease of an individual's response to his or her own cells .- In immune prophylaxis, stress proteins are administered to prevent or reduce the effects in an individual of a pathogen, which can be any virus, microorganism or other organism or substance (e.g., a toxin or toxoid) which causes disease. In preventing or reducing adverse effects of 25 nonviral pathogens (e.g., bacteria, mycobacterial) according to the method of the present invention, an individual's immune response to the nonviral pathogen's stress protein(s) is induced or enhanced through the administration of a vaccine which includes the pathogen's 30 stress protein(s) and, generally, an adjuvant.

Preventing or reducing adverse effects of viral pathogens, as well as preventing cell transformation or reducing the extent to which it occurs, according to the present method, is effected by transiently enhancing an individual's immune surveillance system. In this instance, the causative pathogens (i.e., virus; transforming agent) do not have stress proteins of their own. Enhancement of immune response can be effected by modulating the immune cells by stimulation with a nonviral stress protein (e.g., a bacterial stress protein) or modulating the individual's stress response by any means (e.g., local application of heat).

In immune therapy, such as is used in treating autoimmune diseases, stress proteins known to be involved in the autoimmune response are administered to turn down an individual's immune response by tolerizing the individual to the stress proteins. Alternatively, the immune response to stress protein, which is known to occur in autoimmune disease, is reduced by interfering with the ability of immune cells which respond to stress proteins to do so.

A selected stress protein of the present invention can be administered to an individual, according to the method of the present invention, and result in an immune response which provides protection against subsequent infection by a nonviral organism (e.g., bacteria, mycobacteria, other infectious agents which produce stress proteins). Alternatively, a selected stress protein can be administered to an individual, generally over time, to induce immune tolerance against the selected stress protein. For example, a selected stress

protein can be administered in multiple doses over time in order to induce immune tolerance against an autoimmune disease such as rheumatoid arthritis.

#### Brief Description of the Drawings

Figure 1 is a graphic representation of the homologies between mycobacterial antigens and known stress proteins. Figure 1A is a representation of sequence similarity between portions of the M. tuberculosis 71-kDa antigen (residues 1-204; TB 71kDa) and the E. coli DnaK protein (residues 430-469). Figure 1B is a representation of sequence similarity between portions of the M. tuberculosis 65-kDa antigen (residues 1-540; TB 65kDa) and the E. coli GroEL protein (residues 1-547).

Figure 2 is a comparison of the amino acid sequence of the human P1 protein (573 residues) and the amino acid sequence of the groEL protein (547 residues).

Figure 3 is a comparison of the amino acid sequence of the human Pl protein (573 residues), which is a homolog of groEL protein, and the amino acid sequence of the 65kDa M. leprae protein (540 residues).

Figure 4 is a comparison of the amino acid sequence of the human Pl protein (573 residues), which is a homolog of the groEL protein, and the amino acid sequence of the 65kDa M. tuberculosis protein (540 residues).

## 25 <u>Detailed Description of the Invention</u>

riven saudo a militar quationalita

The present invention is based on the observation that stress proteins are among the major antigens available for presentation to T lymphocytes and may be common immune targets in a broad spectrum of infectious

Immune responses to stress proteins are involved in immune surveillance by the body and a variety of different T cell types has been shown to recognize highly conserved stress protein determinants. observations, described below, suggest a model of immune 05 surveillance in which self-reactive T cells provide a first line of defense against infection or other invasion by pathogens and against cell transformation by recognizing and helping to eliminate stressed autologous cells, as well as cells infected with intracellular bacteria. Without wishing to be bound by this model, it is presented as one means by which it is possible to explain why prokaryotic and eukaryotic cells respond to a variety of potentially damaging stimuli, such as elevated temperature, by increasing the synthesis of a family of 15 proteins, referred to as stress proteins, which are among the most highly conserved and abundant proteins found in nature. 

Investigation of antigens involved in the immune response to the tuberculosis and leprosy bacilli (M. tuberculosis and M. leprae) initially led to the observation that a variety of stress proteins are among the major targets of the immune response, as is described at greater length below.

Further assessment has demonstrated that stress proteins may be common immune targets in a broad spectrum of infectious diseases. Sequence analysis has revealed 70-kDa heat shock protein homologues among major antigens of the protozoan parasites <u>Plasmodium falciparum</u> (Bianco, 30 A.E. et al., Proc. Natl. Acad. Sci. USA 83:8713 9717

A.E. et al., Proc. Natl. Acad. Sci., USA, 83:8713-8717 (1986)) and Schistosoma mansoni (Hedstrom, R. et al., J.

Exp. Med., 165:1430-1435 (1987)) and the malarial parasite Brugia malayi (Selkirk, M.E. et al., J. Cell Biochem., 12D:290 (1988)). Similarly, homologues of GroEL have been found among antigens involved in the immune response to Salmonella typhimurium and Coxiella. 05 Vodkin, M.H. and J.C. Williams, J. Bacteriol, 170:1227 The presence of stress proteins among major immune targets in a variety of human pathogens is support for the idea that the stress response may be a general 10 component of infection and that stress proteins should be considered among candidates for subunit vaccines. All organisms respond to heat by inducing synthesis of heat shock proteins (hsps), which are a group of proteins. This response is the most highly conserved genetic system known and has been shown to occur in every organism, 15 including microorganisms, plants and animals. investigated to date. Many of the characteristics of the response are common to all organisms and the hsps are among the most highly conserved proteins known. For 20 example, hsp90 family and hsp70 family proteins are present in widely diverse organisms. The proteins in each family--even in such diverse organisms--show approximately 50% identity at the amino acid level and at the nonidentical residues, exhibit many similarities 25. Several of the proteins induced by heat are also induced by a variety of other stresses. The hsps or a closely related/similar protein are present in all organisms at normal temperatures and have been shown to have key functions in normal cell metabolism. Lindquist, S. and 30 E.A. Craig, Ann. Rev. Genet., 22:631-677 (1988). the stress response is common to prokaryotes and

eukaryotes and stress proteins are among the most highly conserved in sequence, it is reasonable to expect that an antigen from one pathogen could immunize against another pathogen. Exposure to foreign stress proteins early in life might, in fact, induce a degree of immunity to a variety of infectious agents. If so, this could provide an explanation for the observation that, for many pathogens, only a fraction of infected individuals actually acquire clinical disease.

The following is a description of the relationship which has been observed between stress proteins and the immune response to mycobacterial infection; of the observation and supporting information that stress proteins are immune targets in many non-viral infections; of recognition of the fact that immune responses to conserved stress protein determinants may play an important role in autoimmune pathology in rheumatoid arthritis, as well as in adjuvant arthritis; and of the role of stress proteins in immune surveillance, as well as a model proposed for immune surveillance in which self-reactive T cells provide a first line of defense against infection and cell transformation.

Mycobacterial Stress Proteins are Targets of the Immune
Response

· 1962、 我说的话: 1962年 1000年 1200年 120

An intriguing relationship between stress proteins and the immune response to mycobacterial infection has been observed. A more detailed examination of stress protein determinants and immune response mechanisms is essential to understanding the relationships among stress proteins, infection, and immunity.

In view of the involvement of proteins of M. tuberculosis and M. leprae in humoral and cell-mediated immune responses and to establish the functions of the these proteins in the mycobacterial cell, the DNA 05 encoding several of the M. tuberculosis and M. leprae antigens have been sequenced. It has been demonstrated, as a result, that many of these mycobacterial protein antigens exhibit striking sequence similarity to known stress-induced proteins. Three of the M. leprae and 10 two of the M. tuberculosis protein antigens studied have been shown to exhibit striking sequence similarity to known stress proteins. For reasons discussed in the Exemplification, it is concluded that two of the M. <u>leprae</u> and two of the <u>M. tuberculosis</u> antigens are 15 homologues of the E. coli DnaK and GroEL proteins.

In experimental mice, immunization with mycobacterial lysates elicits antibody responses to at least six M. tuberculosis protein antigens and a similar number of M. leprae protein antigens. Monoclonal anti-20 bodies specific for these proteins have been used to isolate clones from  $\lambda gtll$  DNA expression libraries of  $\underline{M}$ . tuberculosis, and M. leprae. The sequence of the DNA clones revealed that mycobacterial hsp70 (alias 70 kDa antigen) and hsp60 (alias 65 kDa antigen, -groEL) were the major targets of the murine antibody response to both M. tuberculosis and M. leprae. Two additional hsp's, an 18 kDa member of the small hsp family and a 12 kDa homologue of groES, were found among the M. leprae and M. tuberculosis antigens. Young, D.B., et al., Proc. Natl. <u>Acad. Sci., USA</u>, <u>85</u>:4267-4270 (1988); Shinnick, T.M., <u>ec</u> 30 al., Nuc. Acids Res., 17:1254 (1989).

10

The mycobacterial stress proteins are among the immunodominant targets of both murine antibody and T cell In one study which summarized results obtained from 10 laboratories, a collection of 24 murine monoclonal antibodies recognized 6 M. leprae proteins; 7 of these antibodies are directed against 6 different determinants in the M. leprae hsp60. Engers, H.D., et <u>al.</u>, <u>Infect. Immun.</u>, <u>48</u>:603-605 (1985); Mehra, V., <u>et</u> al., Proc. Natl. Acad. Sci., USA, 83:7013-7017 (1986). In a similar study, 3 of 33 monoclonal antibodies raised

against M. tuberculosis recognized the M. tuberculosis hsp60 protein. Engers, H.D., et al., Infect. Immun., 51:718-720 (1986). Finally, limiting dilution analysis indicates that 20% of the mycobacterial-reactive CD4 $^+$  T lymphocytes in mice immunized with M. tuberculosis 15 recognize this antigen. Kaufmann, S.H., et al., Eur. J.

<u>Immunol.</u>, <u>17</u>:351-357 (1987).

Although a rigorous quantitative analysis of the human immune response to mycobacterial stress proteins has not yet been reported, mycobacterial stress proteins 20 are recognized by human antibodies and T lymphocytes and the evidence suggests that these proteins are among the major targets of the human cell mediated immune response. Emmrich, F., et al., J. Exp. Med., 1163:1024-1029 (1985); 25 Mustafa, A.S., et al., Nature (London), 1319:63-66 (1986); Oftung, F., et al., J. Immunol., 138:927-931 (1987); Lamb, J.R., et al., EMBO J., 6:1245-1249 (1987). lymphocytes from patients with mycobacterial infection or from volunteers immunized with mycobacteria have been 30 cloned and tested for their ability to recognize the mycobacterial stress proteins. In each of these studies,

some fraction of the human T cell clones were shown to recognize one or more of the mycobacterial stress proteins.

# Stress Proteins are Immune Targets in Non-viral 1 Infections

The observation that stress proteins are important targets of the immune response to mycobacterial infection and the knowledge that the major stress proteins are conserved and abundant in other organisms suggested that 10 stress proteins are likely to be immune targets in many non-viral infections. Indeed, that is now clearly the case. Antigens from a wide variety of infectious agents have been identified as members of stress protein families. The major stress protein antigen recognized by antibodies in bacterial infections is hsp60. "Common 15 antigen", an immunodominant protein antigen long known to be shared by most bacterial species, turns out to be Shinnick, T.M., et al., Infect. Immun., 56:446 (1988); Thole, J.E.R., et al., Microbial Pathogenesis,  $\underline{4}$ :71-83 (1988). Stress proteins have also been 20 identified as immune targets in most major human parasite infections. Bianco, A.E., et al., Proc. Natl. Acad Sci., USA, 83;8713 (1986); Nene, V., et al., Mol. Biochem. Parasitol., 21:179 (1986); Ardeshir, F., e EMBO J., 6:493 (1987); Hedstrom, R., et al., J. Exp. 25 Med., 165:1430 (1987); Selkirk, M.E., et al., J. Cell Biochem., 12D:290 (1988); Engman, D.M., et al., J. Cell Biochem., 12D: Supplement, 290 (1988); Smith, D.F., et

<u>al.</u>, <u>J. Cell Biochem.</u>, <u>12D</u>:296 (1988). Antibodies to

hsp70 have been identified in the sera of patients

suffering from malaria, trypanosomiasis, leishmaniasis, shistosomiasis and filariasis. Hsp90 is also a target of antibodies in trypanosomiasis and a member of the small hsp family is recognized in some patients with shistosomiasis.

Stress Proteins and Autoimmune Processes Rheumatoid arthritis is characterized by a chronic proliferative and inflammatory reaction in synovial membranes which is thought to involve autoimmune processes. Rat adjuvant arthritis resembles human 10 rheumatoid arthritis in many respects, and has been used as an experimental animal model for human disease. Pearson, C.M., <u>Arthritis Rheum.</u>, 7:80-86 ((1964). Adjuvant arthritis can be induced in rats with a single 15 intradermal injection of killed M. tuberculosis in complete Freund's adjuvant. An autoimmune process involving T lymphocytes appears to be responsible for the generation of the disease. Holoshitz, J., et al., Science, 219:56-58 (1983). T cell lines isolated from

the draining lymph nodes of arthritic rats and propagated in vitro by stimulation with M. tuberculosis-pulsed syngeneic antigen presenting cells can cause a transient form of the disease when transferred to irradiated rats.

Since care was taken in these experiments to exclude the transfer of contaminating M. tuberculosis this results.

25 transfer of contaminating M. tuberculosis, this result strongly suggests that the clinical effects of the disease are a consequence of an autoimmune reaction in which the autoantigen is shared with M. tuberculosis.

The rat and  $\underline{M}$ . tuberculosis antigens recognized by 30 the arthritogenic T cells have been sought for a number

of years. A number of different proteins present in synovial membranes have been proposed to be the cross-reactive rat antigen, but were later discounted as procedures for the purification of these proteins improved. van Eden, W., et al., Proc. Natl. Acad. Sci., 05 <u>USA</u>, <u>82</u>:5117-5120 (1985); Holoshitz, J., <u>et al.</u>, <u>Science</u>, 219:56-58 (1983). The M. tuberculosis antigen recognized by the arthritogenic T cells was recently shown to be a 65 kDa protein (van Eden, W., et al., Nature, 331:171 (1988), which has now been shown to be hsp60 (see the 10 Exemplification). Using a combination of truncated recombinant 65 kDa proteins and peptides, a nine amino acid epitope of hsp60 has been identified as the minimum stimulatory sequence for arthritogenic T cell clones in proliferation assays. Now that it is clear that some 15 arthritogenic T cells recognize the mycobacterial hsp60, it is quite possible that the rat autoantigen is also hsp60. ...

The results obtained in the adjuvant arthritis model

led investigators to determine whether T lymphocytes from human rheumatoid arthritis patients also recognize mycobacterial antigens. These investigators have found not only that patients with rheumatoid arthritis have T cells that recognize M. tuberculosis antigens, but that these T cells have diverse phenotypes. Substantial proliferative responses to mycobacterial extracts are observed with uncloned T cells (predominantly CD4<sup>+</sup>) from both synovial infiltrates and peripheral blood, although responses are generally greater in synovial infiltrates.

Abrahamson, T.G., et al., Scand. J. Immunol., 7:81-90 (1978); Holoshitz, J., et al., Lancet ii, 305-306 (1986).

Holoshitz et al. found that 4 of 5 T cell clones isolated from human rheumatoid synovia which respond to M. tuberculosis antigens were CD4 CD8 cells with  $\gamma/\delta$  T cell receptors. Holoshitz, J., et al., Nature, 05 339:226-229 (1989). This observation is interesting because  $\gamma/\delta$  T cells have yet to be assigned a role in immunity. One of the  $\gamma/\delta$  clones was tested for its ability to respond to purified mycobacterial hsp60 and was found to be positive in proliferation assays. Due to 10 the conserved nature of stress proteins, these T cells have the potential for autoreactivity. Lamb and it is a second coworkers have shown that polyclonal T cells from synovial infiltrates recognize both mycobacterial hsp60 and hsp70. Lamb, J.R., et al., Intl. Immunol., in press (1989). The population of T cells that recognize the 15 mycobacterial stress proteins were shown to respond to E. coli hsp60 and hsp70 and, most interestingly, human hsp70 purified from heat shocked macrophages. Thus, immune responses to conserved stress protein determinants, 20 perhaps initiated by bacterial infection (not necessarily by mycobacteria), may play an important role in autoimmune pathology in rheumatoid arthritis, as well as in adjuvant arthritis. solis the proprise M. <u>Interouseis</u> antipers, solisses

A variety of different T cell types has now been shown to recognize highly conserved stress protein determinants. The ability of cells to respond to stress by increasing the levels of the highly conserved stress proteins; the presence of T cells of diverse phenotypes in healthy individuals that are capable of recognizing

self stress protein determinants; and observations that

stress responses are induced by viral infection and by cell transformation, all suggest a model of immune surveillance in which self-reactive T cells provide a 05 first line of defense against infection and transformation by recognizing and helping to eliminate stressed autologous cells, as well as cells infected with intracellular bacteria. The pool of lymphocytes that recognize conserved stress protein determinants might be 10 induced during establishment of natural microbial flora on the skin and in the gut, and maintained by frequent stimulation by bacteria and viruses as well as other stressful stimuli encountered during a normal lifetime. This model is attractive because it provides a way in which the immune system could exploit the existence of 15 conserved epitopes in stress proteins to respond immediately to antigenically diverse pathogens and cellular changes, producing an initial defense that need not await the development of immunity to novel antigens, 20 Stress protein induction occurs in eukaryotic cells following infection by diverse viruses in vitro. Collins, P.L., and Hightower, L.E., J. Virol., 44:703-707 (1982); Nevins, J.R., Cell; 29:913-939 (1982); Garry R.F., et al., Virology, 129:391-332 (1988); Khandjian, E.W. and Turler, H., Mol. Cell Biol ? 3:1-8 (1983) LaThangue, N.B., et al., EMBO J., 3:267-277 (1984). CTL that recognize these neo-antigens could limit the spread of virus by killing infected cells, possibly before substantial amounts of mature virus are assembled, and by 30 secreting the lymphokine  $\gamma$ -interferon. Pestka, S., in: Methods Enzymol., Interferons, Part A., Vol. 79, Academic

Press, New York, pp.667 (1981). Evidence consistent with this idea is emerging. Koga et al., have shown that infection of primary murine macrophages with CMV rendered them susceptible as targets for MHCOI restricted CD8<sup>+</sup> CTL specific for linear epitopes of M. tuberculosis hsp60. Koga, T., et al. Although the epitope recognized by these CTL on infected macrophages was not defined, it is tempting to speculate that a cross-reactivity with self hsp60 epitopes is being observed. Indeed, the same groups showed that a homologous hsp60 is constitutively present in macrophages and is upregulated by γ-interferon stimulation.

T cells capable of recognizing autologous cells stressed by transformation could help eliminate nascent tumour cells. Stress proteins appear to be produced at 15 high levels in at least some transformed cells. Bensaude, O. and Morange, M., EMBO J., 2:173-177 (1983). An 86 kDA murine tumour antigen has been found to be homologous to representatives of the hsp90 family in 20 yeast and <u>Drosophila</u>. Ullrich, S.J., <u>Proc. Natl. Acad.</u> Sci., USA, 83:3121-3125 (1986). Immunization of mice with the purified protein led to inhibition of tumor growth in 95% of experimental animals that had been seeded with cultured tumor cells. All of the protected mice had high titres of anti-hsp90 serum antibody which 25 was able to precipitate murine hsp90 from lysates of heat shocked mouse embryo cells. Again, a role for autoreactive lymphocytes is implied.

The lymphocytes which recognize conserved stress
protein determinants must be capable of discriminating between normal and stressed cells. Since many stress

proteins are constitutively expressed in normal cells, although at lower levels than in stressed cells, the potential for autoreactivity is ever-present. Normal cells may escape destruction by expressing only substimulatory levels of stress protein determinants on 05 their surfaces. In addition, stress proteins may only be processed and presented during stress, and it may be relevant that many stress proteins have altered intracellular locations during stress. Finally, immune regulatory networks may prevent activation of 10 autoreactive T cells under normal conditions. The regulatory constraints required by this system might occasionally break down, perhaps during stress caused by bacterial or viral infections, leading to autoimmune 15 disease. Rhematoid arthritis may be such a disease. Modulation of Immune Response

The precise relationship between stress proteins and the host immune response to infection is as yet undefined. When cells are subjected to a variety of 20 stresses, they respond by selectively increasing the synthesis of a limited set of stress proteins. Some stress proteins, including the products of dnaK and groEL, are major constituents of the cell under normal growth conditions, and are induced to even higher levels 25 during stress. Lindquist, S., Annu. Rev. Biochem. 1151-1191 (1986); Neidhardt, F.C. and R.A. VanBogelen, In Escherichia coli and Salmonella Typhimurium, Cellular and Molecular Biology, (eds. Neidhardt, F.C., Ingraham, J.L. Low, K.B. Magasanik, B. Schaechter, M. and Umbarger, 30 H.E.) Am: Soc. Microbiol., Washington, D.C., pp. 1134-

1345 (1987). It has now been demonstrated that stress-

related proteins are targets of the immune response. It is reasonable to expect that immunodominant antigens would be found among such abundant proteins, as has been shown to be the case.

Of Stress-induced proteins or their functional equivalents, can be used to immunize an animal against a nonviral infection or, alternatively, other selected stress-induced proteins, or their functional equivalents, can be used to induce immune tolerance in an animal.

- According to the method of the present invention, it is possible to modulate the immune response in an individual by altering the individual's response to stress proteins. In particular, it is possible to modulate an individual's response to a pathogen (e.g., bacterium, mycobacterium,
- virus, or other organism or agent, such as parasites, toxins, toxoids, which causes disease or cell transformation); and to modulate an individual's autoimmune response, such as occurs in some forms of arthritis.
- It is possible to use stress proteins, as described herein, as a vaccine which, after administration to an individual, produces or enhances an immune response in that individual, providing protection against subsequent infection. Because, as demonstrated, these are proteins in which there are regions of highly conserved amino acid sequences and such proteins have been shown to be ubiquitous targets in mycobacterial, bacterial and other infections, they can be used to elicit an equally

For example, in the case of a pathogen which 30 expresses stress proteins, such as nonviral pathogens, two approaches to enhancing an individual's immune

ubiquitous immune response.

response (and, thus, reducing the pathogen's effects) can be used.

First, because the nonviral pathogen's stress proteins are distinguishable from those of the host, it 05 is possible to induce an immunoprophylactic response specific to the pathogen's stress proteins. This can be carried out by administering a vaccine which includes all or a portion (e.g., sufficient sequence to have the desired stimulatory effect on immune response) of the pathogen's stress protein or of another protein having an amino acid sequence sufficiently similar to that of the stress protein sequence to stimulate the immune response to the pathogen. Alternatively, highly conserved stress protein determinants, such as those shown to be 15 recognized by a variety of T cells, can be administered as a type of "general" vaccine. In either case, the immune response to the stress protein sequence will be increased and effects of the nonviral pathogen will be reduced (decreased, prevented or eliminated).

Second, it is also possible to induce or enhance the immune surveillance system or response which is directed to stressed host cells. This is described further in the context of enhancing immune response in those instances in which the pathogen (e.g., a virus, transforming agent) does not have (express) its own stress proteins (i.e., stress proteins distinguishable from host stress proteins).

20

25

30

The vaccine administered to induce or enhance immune response to nonviral pathogens includes a stress protein of the pathogen against which an immune response is desired, a portion of that protein of sufficient size to

stimulate the desired immune response or a protein or amino acid sequence (e.g., a polypeptide) which is the functional equivalent of the pathogen stress protein in that it is sufficiently homologous in sequence to that of 05 the stress protein to be capable of eliciting the desired response. The term "sufficiently homologous in sequence to that of the stress protein" means that the sequence of the protein or polypeptide will generally show approximately 50% identity with the stress protein amino acid sequence. The vaccine can also include an adjuvant, 10 an appropriate carrier and an appropriate buffer. protein or amino acid sequence present in the vaccine can be produced using known techniques. For example, it can be obtained (isolated) from a source in which it occurs 15 in nature, can be produced by cloning and expressing the gene encoding the desired stress protein or stress protein portion or can be synthesized chemically or mechanically. 

In the case of a pathogen, such as a virus or a 20 transforming agent (i.e., an agent whose activity results in production or formation of cancer cells), which does not express its own stress proteins, the following approach is used to enhance immune response or normal immune surveillance (i.e., the ability of the immune system to recognize self as well as foreign proteins). 25 A vaccine which includes a bacterial, mycobacterial or other stress protein can be administered. Although there are no viral stress proteins to be used for this purpose, administration of such a vaccine will enhance the existing immune surveillance system. In addition, immune 30 surveillance can also be enhanced by applying local heat

or any other substances or changes in condition which induce the stress response in the individual being treated. (This can also be employed in conjunction with the vaccine, described previously, administered to enhance immune response to a stress protein-producing pathogen.) It is known that increased levels of stress proteins are produced in many types of cancer cells. Enhancement of the immune surveillance system, as described, can be used to facilitate destruction and/or to prevent progression or establishment of cancer cells.

The method of the present invention can also be used to modify or modulate an individual's response to his or her own cells (e.g., as in autoimmune diseases). are at least two ways in which the present invention can be used immunotherapeutically. First, stress proteins, 15 such as heat shock protein (hsp) 70 and hsp60, are known to be involved in autoimmune disease. It is, thus, possible to turn down an individual's response to "self" by administering the appropriate stress protein(s) in such a manner that the individual becomes more tolerant 20 of the protein. Second, because it is known that the immune response in autoimmune diseases is to stress proteins, it is possible to selectively inhibit or interfere with the ability of immune cells which normally interact with such proteins to do so. This can be done for example, by administering monoclonal antibodies that bind to specific T cell receptors and delete or disable such cells. Alternatively, rather than knocking out immune cells, the stress response in all cells can be turned down by administering a drug capable of reducing a

cell's-ability to undergo the stress response.

example, a drug targeted to or specific for heat shock transcription factor, which is needed to stimulate heat shock genes, can be administered. The transcription factor is rendered nonfunctional or subfunctional and, as a result, cells' ability to undergo the stress response is also lessened.

The present invention is further illustrated by the following exemplification, which is not intended to be limiting in any way.

5.148

### 10 <u>EXEMPLIFICATION</u>

Recombinant DNA Clones. The isolation and characterization of M. tuberculosis and M. leprae  $\lambda$ gtll genomic DNA clones with murine monoclonal antibodies have been described. Husson, R.N. and Young, R.A., Proc.

- Natl. Acad. Sci., USA, 84:1679-1683 (1987); Young, R.A., et al., Nature, (London) 316:450-452 (1985). DNA was isolated from these clones and was manipulated by standard procedures. Davis, R.W., Advanced Bacterial Genetics: A Manual for Genetic Engineering (Cold Spring Harbor Lab., Cold Spring Harbor, NY), (1980).
- DNA Sequence Analysis and DNA was subcloned into vector M13mp18 or M13mp19 (New England Biolabs), as suggested by the supplier. Dideoxynucleotide chain termination reactions and gel electrophoresis of the sequenced produced were as described. Davis, R.W., Advanced

Bacterial Genetics: A Manual for Genetic Engineering (Cold Spring Harbor Lab., Cold Spring Harbor, NY),

(1980). DNA sequences were determined for both strands of DNA. Computer analysis of sequences with UWGCG programs was as described by Devereux, J., et al. Nucleic Acids Res., 12:387-395 (1984).

- O5 Immunoblot Analysis. Escherichia coil strain TG1 was transformed with the following plasmids by standard procedures (Maniatis, T., et al., Molecular Cloning, A Laboratory Manual (Cold Spring Harbor Lab., Cold Spring Harbor, NY) (1982), with selection for ampiclling the content of the con
- resistance: pND5, a derivatrive of pBR325 containing the E. coli groE genes (Jenkins, A.J., et al., Mol. Gen. Genet, 202:446-454 (1986); pUC8 (Vic?, J., Gene, 19:259-268 (1982); pUC8 with insert DNA for λgtll clone Y3178 (M. leprae 65-kDa antigen, Young, R.A., et al.,
- Nature, (London) 316:450-452 (1985)) ligated in the EcoRI site.

Overnight cultures of <u>E. coli</u> strains in Luria-Bertani (LB) medium were centrifuged and resuspended in isotonic phosphate-buffered saline at a cell density corresponding to an absorbance of 20 at 60 nm. An equal volume of sample buffer containing 2% (wt/vol) polycrylamide gels in the presence of NaDodSO<sub>4</sub> was added, and, after heating on a boiling water bath for 2 min, 5-ml samples were electrophoresed on 12% (wt/vol)

- polycrylamide gels in the presence of NaDodSO<sub>4</sub>. Blots were prepared by electrophoretic transfer of the proteins to a nitrocellulose membrane, and binding of monoclonal antibodies was assayed with a peroxidase-conjugated secondary antibody as described. Young., D.B.. et al.,
- 30 <u>Infect. Immun.</u>, <u>55</u>:1421-1425 (1987).

Six M. tuberculosis and six leprae proteins have been implicated in the immune response to the mycobacterial pathogens (T1). To obtain clues to the normal cellular function of several of these 05 mycobacterial antigens, DNA clones encoding these proteins, isolated by using monoclonal antibodies to probe lambdagtll libraries (Husson, R.N. and Young, R.A., Proc. Natl. Acad. Sci., USA, 84:1679-1683 (1987); Young, R.A., et al., Nature, (London) 316:450-452 (1985)) were 10 subjected to sequence analysis. The sequences elucidated have been submitted to the GenBank sequence database.

The Mycobacterial 71-k Da Antigen. The 71-k Da antigen of M. tuberculosis is recognized by human T cells during infection (Table).

្នាល់ ស្ត្រី ដែលប្រឹត្តិបន្ទេស នេះការសេស ស. ៤.១៤, ខេត្ត នៃនៅនេះ ប្រែក្រសួន ស. និ. ស.ស. ៤

នៃ ប្រទេទី សក្សាទីទី **នូក្សាដ៏ដែលវី នំ** ក្រ ប្រួលវិ ខ្លួក ខ្<mark>នួលវិទ្</mark>ទី ប្រែក្រុង ស្រុខ ខេត្ត ភេទស្រ

ကြောင်းသည်။ နေရိုင်းကြုံရှိနေရာန် ရောက်သည်။ မေရိုင်းကြုံရှိနှင့် မြောက်သည်။ မြောက်သည်။ မေရိုင်းများကြုံရှိနှင့်

	TABLE	Mycobacteri	al protein ant	tigens	1.
4	·••				
		Recogn	ized Subject	ed Homology	·
		by huma	an T to seque	ence with kno	wn
	<u>Protein,</u>	kDA cells	analysis	<u>proteins</u>	
05	M. tubero	1			
03		culosis			
	71	<b>+</b>	+ -	DnaK	• • •
	65*	• • • • • • • • • • • • • • • • • • •	• • • • • • • • • • • • • • • • • • •	GroEL	
	38	<b>+</b>	•	•	
•	19	<b>+</b>	+ .	None	
10	14		-		
	12.	ND	<u></u> -	•	
				· .	•
	M. leprae			• • • • • • • • • • • • • • • • • • • •	
	. 70 .	ND .		DnaK	
	6.5		······································	GroEL	
15	36	- A+			
	28	+	- ·		
	18	- ( +		Plant Hs	
	12	ND	· · · -	•	
	•				•

Mycobacterial protein antigens, their recognition by human T cells, and homology of the deduced mycobacterial protein sequences to known proteins are summarized.

ND, not determined; +, yes; -, no

unpublished data.

<sup>\*</sup> Includes data derived from study of the 65-kDA

25 antigens of M. bovis BCG (bacillus Calmette-Guerin),
which is identical to the M. tuberculosis 65-kDA antigen.

+ A. S. Mustafa, J. R. Lamb, D. Young and R. A. Young,

The insert DNA of lambdagtll clone Y3271 (Husson, R.N. et al., Proc. Natl. Acad. Sci., USA, 84:1679-1683 (1987), was sequenced to obtain amino acid sequence information for the 71-kDa antigen of M. tuberculosis. This clone produces a beta-galactosidase fusion protein containing 0.5 the carboxyl-terminal one-third of the 71-kDa antigen exhibits 40% amino acid sequence identity with the comparable segment of the  $\underline{\mathtt{dnaK}}$  gene product from  $\underline{\mathtt{E}}$   $\underline{\mathtt{coli}}$ (Bardwell, J.C., et al., Proc. Natl. Sci., USA, 10 81:848-852 (1984)), (Fig. 1). Fig 2A shows the extent of sequence similarity between portions of the mycobacterial and the E. coli 70-k Da polypeptides. Sequences transcriptionally downstream from the mycobacterial 71-k Da gene predict a 356-amino acid protein homologous to the E. coli dnaJ gene product (unpublished data), 15 indicating that the E. coli dnaK-dnaJ operon structure is conserved in  $\underline{M}$ . tuberculosis and consistent with the conclusion that the mycobacterial 71-kDa antigen is a homologue of the E. coli dnaK gene product. The product of the  $\underline{dnak}$  gene is a member of the 70-kDa heat shock 20 protein family that is highly conserved among prokaryotes and eukaryotes (Bardwell, J.C., et al., Proc. Natl. Sci., <u>USA</u>, <u>81</u>:848-852 (1984); Lindquist, S., <u>Annu. Rev.</u> Biochem., 55:1151-1191 (1986). Therotty isizerouse with

The M. leprae 70-k Da antigen cross-reacts with monoclonal antibodies directed to the M. tuberculosis and M. leprae are both members of the 70-k Da heat shock protein family of stress proteins.

10

The Mycobacterial 65-kDa antigen. The 65-kDa antigens of M. tuberculosis and M. leprae are involved in in the human T-cell response to mycobacterial infection (Table). Genes encoding these proteins have been isolated (Husson, R.N. and Young, R.A., Proc. Natl. Acad. Sci., USA, 84:1679-1683 (1987); Young, R.A., et al., Nature, (London) 316:450-452 (1985)) and sequenced (Shinnick, T.M., J. Bacteriol., 169:1080-1088 (1987); Mehram, V., et al., Proc. Natl. Acad. Sci., USA, 83:7013-7017 (1986)), revealing that the amino acid sequences of the 65-kDa antigens of M. tuberculosis and M. leprae are 95% identical. These protein sequences exhibit no significant sequence similarity to proteins in the GenBank database.

Identification of these proteins was based on the observation that some monoclonal antibodies directed against the mycobacterial 65-kDa antigens cross-react with an <u>E. coli</u> protein of 60kDa. <u>E. coli</u> cells transformed with the plasmid pND5 (Sanger, F., et al., Proc. Natl. Acad. Sci. USA. 74:5463-5467 (1977) which

- Proc. Natl. Acad. Sci., USA, 74:5463-5467 (1977), which contains the E. coli gro E genes, had been shown to accumulate large amounts of the 60-kDa protein. A comparison of the mycobacterial 65-kDa protein sequences with those determined for E. coli groEL (C. Woolford, K.
- 25 Tilly, C. Georgopoulous, and R.H., unpublished data) revealed the extent of the sequence similarity as shown in Fig. 1B.

The 60-kDa Gro EL protein is a major stress protein in  $\underline{E.\ coli}$ . Lindquist, S.,  $\underline{Annual\ Rev.\ Biochem.}$ ,

30 <u>55</u>:1151-1191 (1986); <u>Nature</u>, <u>333</u>:330-334 (1988). There is some evidence that the mycobacterial 65-kDa proteins

accumulate in response to stress: Mycobacterium bovis
BCG (bacillus Calmette-Guerin) cultures grown in
zinc-deficient medium are substantially enriched in this
protein (De Bruyn, J., et al., Infect. Immun., 55:245-252
(1987)). This infers that the 65-kDa proteins of M.
tuberculosis and M. leprae are homologues of the E. coli
Gro EL protein.

Other Mycobacterial Antigens. T lymphocytes that respond to the M. tuberculosis 19-kDa antigen and the M.

leprae 18-kDa antigen have been observed in humans with tuberculosis and leprosy, respectively (Table 1). DNA encoding these antigens was sequenced from the Agtll clones Y3148 (Husson, R.N. and Young, R.A., Proc. Natl. Acad. Sci., USA, 84:1679-1683 (1987); and Y3179 (Young, R.A., et al., Nature, (London) 316:450-452 (1985)), respectively. The M. tuberculosis 19-kDa protein sequence predicted from the DNA exhibited no significant

However, the M. leprae 18-kDa protein sequence was

similar to the soybean 17-kDa heat shock protein, a protein representation of a major class of plant heat shock proteins (Schoffl, F. and Van Bogelen, R.A., In:

Escherichia coli and Salmonella typhimurium, Cellular and Molecular Biology, Am. Soc. Microbiol., Washington, D.C.

(1987).

#### 

Those skilled in the art will recognize, or be able to ascertain using no more than routine experimentation, many equivalents to the specific embodiments of the invention described specifically herein. Such equivalents are intended to be encompassed in the scope of the following claims.


#### CLAIMS


- 1. A vaccine comprising all or a portion of a selected stress protein or all or a portion of a protein having an amino acid sequence sufficiently homologous to the amino acid sequence of the stress protein.
- A vaccine of Claim 1 in which the stress protein is a mycobacterial stress protein or a protein having an amino acid sequence sufficiently homologous to the amino acid sequence of the mycobacterial stress protein.
  - 3. A composition for use as an agent to induce immune tolerance, comprising a selected stress protein.
- 4. A composition for use in treating an autoimmune disease, comprising all or a portion of a selected stress protein or all or a portion of a protein having an amino acid sequence sufficiently homologous to the amino acid sequence of the stress protein.
- 20 5. A composition of Claim 4 for treating rheumatoid arthritis.

a afoniosas ens discade

6. A vaccine for use in enhancing in an individual the immune response to a nonviral pathogen, comprising all or a portion of a stress protein of the nonviral pathogen against which the enhanced response is desired.

- A vaccine of Claim 6 in which the stress protein is a mycobacterial stress protein.
- 8. A method of producing or enhancing an immune response in an animal, comprising administering to the animal a selected stress protein, in sufficient quantity to elicit the desired immune response.
- A method of immunizing an animal against a subsequent nonviral infection, comprising administering to the animal a selected stress protein, in sufficient quantity to produce an immune response.
- 10. A method of inducing in an individual immune tolerance against a protein, comprising administering to the individual all or a portion of a selected stress protein or all or a portion of a protein having an amino acid sequence sufficiently homologous to the amino acid sequence of the stress protein, under conditions appropriate for induction of the desired tolerance.
- 20 11. A method of Claim 10, wherein the protein is a protein associated with rheumatoid arthritis.








FIGURE 2

	⊷ 、	10	20	30	40	50	09	. 70
HUMP1 GROEL		PTVFRQMRP	VSRVLAPHL:	TRAYAKDVKF( :::::	SADARALMLO	GVDLLADAVA:::::::::::::::::::::::::::::::	MLRLPTVFRQMRPVSRVLAPHLTRAYAKDVKFGADARALMLQGVDLLADAVAVTMGPKGRTVIIEQSWGS : :::::::::::::::::::::::::::::::::::	'IEQSWGS : : LDKSFGA
	71	80	06	100	110	120	130	140
HUMP1 GROEL	PKVT : : PTIT	PKVTKDGVTVAKS::::::PTITKDGVSVARE	IDLKDKYKN] : :: : IEPEDKFENN	i caklvqdvan IIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIII	NNTNEEAGDG1 : :::: KANDAAGDG1	AGDGTTTATVLARS] :::::::::: AGDGTTTATVLAQA]	IDLKDKYKNIGAKLVQDVANNTNEEAGDGTTTATVLARSIAKEGFEKISKGANPVEI : :::::::::::::::::::::::::::::::::::	GANPVEI : :: GMNPMDL
HUMP1 GROEL	141 RRGVN :: KRGIL	150 SVMLAVDAVIA SIDKAVTAAVE	160 ELKKQSKPVT	170 TTPEEIAQVAT :::::	180 .:sangdkeign: :::::::::::::::::::::::::::::::::::	190 :NIISDAMKKV : :: ::	1 150 160 170 180 190 · 200 210  RRGVMLAVDAVIAELKKQSKPVTTPEEIAQVATISANGDKEIGNIISDAMKKVGRKGVITVKDGKTLNDE  :: :: :: :: :: :: :: :: :: :: :: :: ::	210 GKTLNDE : :::
HUMP1 GROEL	211 LEIIE : : LDVVE	220 EGMKFDRGY:::::::EGMQFDRGYI	230 , ISPYFINTSK ::::: LSPYFINKPE	240 GQKCEFQDAY TGAVELESPF	250 VLLSEKKISS :: ::: ILLADKKISN	260 IQSIVPALEI : ::: IREMLPVLEA	1 220 230 240 250 260 270 280, LEITEGMKFDRGYISPYFINTSKGQKCEFQDAYVLLSEKKISSIQSIVPALEIANAHRKPLVIIAEDVDG : :::::::::::::::::::::::::::::::::::	280 IAEDVDG:::::::
HUMP1 GROEL	281 FALST :::: EALAT	290 STLVLNRLKVG : : :	300 SLQVVAVKAP : ::::: IVKVAAVKAP	310 GFGDNRKNQL:::::::	320 KDMAIATGGA : : : : : : : : : : : : : : : : : : :	330 VFGEEGLTLN : :: VISEE-IGME	1 290 300 310 320 330 340 350 EALSTLVLNRLKVGLQVVAVKAPGFGDNRKNQLKDMAIATGGAVFGEEGLTLNLEDVQPHDLGKVGEVIV  ::::::::::::::::::::::::::::::::::	350 KVGEVIV :
	351	360	370	380	390	400	410	420
HUMP1 GROEL	TKDDA::	MLLKGKGDR : : TIIDGVGEE	AQIEKRIQE::::::AAIOGRVAO.	IIEQLDVTTS:::IROOIEEATSI	EYEKEKLNER.::::::	LAKLSDGVAV	TKDDAMLLKGKGDKAQIEKRIQEIIEQLDVTTSEYEKEKLNERLAKLSDGVAVLKVGGTSDVEVNEKKDR :: :: :: :: :: :: :: :: :: :: :: :: ::	NEKKDR::::

...

eastury and a

STATE OF THE STATE ्रक्षा स्वास्त्र स्था

Comparation of いいできまして

TO THE PARTY OF TH

CARRANGE CONTROL OF THE CARRAGE CONTROL OF

.

	1
•	
TINO C	4
Z	2
C	j
Č	í
_	•
C	1
Ĺ	1
α	í
	)
Ü	)
Ĕ	1
FIGHPE	
	•

	421	430	440	450	460	470	480	490
HUMP1	VTDAI	LNATRAAVE	EGIVLGGGCAL	LRCIPALDSL	TPANEDQKI	VTDALNATRAAVEEGIVLGGGCALLRCIPALDSLTPANEDQKIGIEIIKRTLKIPAMTIAKNAGVEGSLI	PAMTIAKNAC	VEGSLI
GROEL	VEDAI	EDALHATRAAVEEG	EGVVAGGGVAL	VAGGGVALIRVASKLADLRGQNEDQ	::: RGQNEDQNV	: ::::::::::::::::::::::::::::::::::::	PLRQIVLNCGEEPSV	SEEPSVV
	491	500	510	520	530	540	550	260
HUMP1	VEKIM	<b>1QSSSEVGYI</b>	DAMAGDFVNMV	EKGIIDPTKV	VRTALLDAA	VEKIMQSSSEVGYDAMAGDFVNMVEKGIIDPTKVVRTALLDAAGVASLLTTAEVVVTEIPKEEKDPGMGA	VVTEIPKEEK	DPGMGA
GROEL	ANTVK	GGDGNYGYNAL		::::: DMGILDPTKV	TRSALQYAA	:: :: :: :: :: :: :: :: :: :: :: :: ::	:: :: MVTDLPKND-	:: AADLGA
	561	570	·					
HUMP1	MGGMG	MGGMGGGMGGGMF	ĨŦ,					. •

score = 4667, 5 breaks identities out of 545 possible matches between residues Total 276

AGGMGGMGGMM-

GROEL

25 random runs Alignment score = 65.34 SD Standard deviation =

Mean = 3429.48

18.94

٠.	•
E C	3
ρ	7
	)
Ľ	•
F.7	
-	,

	<b></b>	10	20	30	40	50	09	70
HUMP1 ML65K	MLRLE : M	TVFRQMRP	MLRLPTVFRQMRPVSRVLAPHLTRAYAKDVKFGADARALMLQGVDLLADAVAVTMGPKGRTVIIEQSWGS: : ::::::::::::::::::::::::::::::::::	LAPHLTRAYAKDVKFGADARALMLQGVDLLADAVAVTMGPKGRTVIIEQSWGS :: ::::::::::::::::::::::::::::::::::	NDARALMLQGV :: EEARRGLERGI	/DLLADAVAV' ::::: .NSLADAVKV'	IMGPKGRTVI : ::::: ILGPKGRNVV	IEQSWGS : :: LEKKWGA
	71	80	06	100	110	120	130	140
HUMPI ML65K	PKVTK : : PTITN	PKVTKDGVTVAKSIDL: ::::::::	PKVTKDGVTVAKSIDLKDKYKNIGAKLVQDVANNTNEEAGDGTTTATVLARSIAKEGFEKISKGANPVEI : :::::::::::::::::::::::::::::::::::	KDKYKNIGAKLVQDVANNTNEEAGDGTTTATVLARSIAKEGFEKISKGANPVEI : : :: :: :: :: :: :: :::::::::::::::	TTNEEAGDGTT::::::::::::::::::::::::::::::::::	TATVLARSII ::::: TATVLAQALI	AKEGFEKISKO ::: /KEGLRNVAAO	SANPVEI :::: SANPLGL
	141	150	160	170	180	190	200	210
HUMP1 ML65K	RRGVM :: KRGIE	LAVDAVIA ::: :KAVDKVTE	RRGVMLAVDAVIAELKKQSKPVTTPEEIAQVATISANGDKEIGNIISDAMKKVGRKGVITVKDGKTLNDE :: :::: :::: ::::::::::::::::::::::::	PEEIAQVATI : : : : :KEQIAATAAI	SANGDKEIGN::::sA-GDQSIGD	IIISDAMKKVG : ::::	SRKGVITVKD( : :::: SNEGVITVEE	SKTLNDE:
	211	220	230	240	250	260	270	280
HUMP1 ML65K	LEIIE	GMKFDRGY:	LEIIEGMKFDRGYISPYFINTSKGQKCEFQDAYVLLSEKKISSIQSIVPALEIANAHRKPLVIIAEDVDG :: ::: :: :: :: :: :: :: :: :: :: :: ::	GKCEFQDAYV : : :QEAVLEEPYI	LLSEKKISSI :: :: LLVSSKVSTV	QSIVPALEIA : :: KDLLPLLEKV	NAHRKPLVI) : : : 71QAGKSLLI)	CAEDVDG
	281	290	300	310	320	330	340	350
HUMP1 ML65K	EALST :::: EALST	LVLNRLKVC::: LVVNKIRG1	EALSTLVLNRLKVGLQVVAVKAPGFGDNRKNQLKDMAIATGGAVFGEEGLTLNLEDVQPHDLGKVGEVIV :::::::::::::::::::::::::::::::::::	FGDNRKNQLK ::::::::	DMAIATGGAV ::::::: DMAILTGAQV	FGEEGLTLNI :::: ISEE-VGLTI	EDVQPHDLG ::: ENTDLSLLG	CVGEVIV
	351	360	370	380	390	400	410	420
нимР1	, TKDDA	MLLKGKGDF	TKDDAMLLKGKGDKAQIEKRIQEIIEQLDVTTSEYEKEKLNERLAKLSDGVAVLKVGGTSDVEVNEKKDR	IEQLDVITSE :	YEKEKLNERL : ::: :::	AKLSDGVAVL	KVGGTSDVEV	NEKKDR
ML65K	TKDET	TIVEGAGD1	tkdettivegagdtdaiagrvaqirteiensdsdydreklqerlaklaggvavikagaatevelkerkhr	RTEIENSDSD	YDREKLQERL	AKLAGGVAVI	KAGAATEVEI	,керкнр

to white her

TO THE STREET OF 以後各種問 情行 は

*े* 03

Little Brother.

Mean =

Standard deviation =

47.73 SD

25 random runs Alignment score

1		١
•	-	
	E	1
1	2	2
i	c	1
(		j
	~	
1	Ŀ	1
1		4
;	Ξ	)
į	2	)
۱	_	į
1	ì	4

149		•					
149	TRAAVEEGIV	VTDALNATRAAVEEGIVLGGGCALLRCIPALDSLTPANEDQKIGIEIIKRTLKIPAMTIAKNAGVEGSLI	IPALDSLTP	ANEDQKIGI	SILKRTLKIPA	MTIAKNAGV	EGSLI
56	::::::: IEDAVRNAKAAVEEGIV	AGGGVTLLQA	APALDKLKL	: rgdeat-gan	::::::::::::::::::::::::::::::::::::::	KQIAFNSGM	: EPGVV
56	500	510	520	530	540	550	560
56	SSEVGYDAMA	GDFVNMVEKG	IIDPTKVVR1	ALLDAAGVA	VEKIMOSSSEVGYDAMAGDFVNMVEKGIIDPTKVVRTALLDAAGVASLLTTAEVVVTEIPKEEKDPGMGA	, TEIPKEEKD)	PGMGA
56	SVGHGLNAAT	: GEYEDLLKAG	:::: VADPVKVTRS	:: :: :: :: DPVKVTRSALQNAASIAC	AEKVRNLSVGHGLNAATGEYEDLLKAGVADPVKVTRSALQNAASIAGLFTT-EAVVADKPEKTAAPASDP	ADKPEKTAAI	PASDP
	570						
	MGGGMF						
ML65K TGGMGG-MD	: MDF				•		
Total score = 4552, 7 breaks 255 identities out of 540	52, 7 breal out of 540	eaks 540 possible matches between residues	matches b	etween re	sidues		

IGURE 4

7.0	QSWGS ::	140	NPVEI :: NPLGL	210 TLNDE	: TFGLQ	280	EDVDG :::: EDVEG	350	GEVIV :: RKVVV	420	SKKDR	SRKHR
60	AYAKDVKFGADARALMLQGVDLLADAVAVTMGPKGRTVIIEQSWGS:::::::::::::::::::::::::::::::::::	130	AKEGFEKISKGA ::: RKEGLRNVAAGA	160 170 180 190 200 210, COSKPVTTPEEIAQVATISANGDKEIGNIISDAMKKVGRKGVITVKDGKTLNDE	SNEGVI TVEESN	270	NAHRKPLVIIA ::: ::: /IGAGKPLLIIA	340	EDVQPHDLGKV	410	KVGGTSDVEVNI	KAGAATEVELKE
20	ODLLADAVAV:::::::::::::::::::::::::::::::	120	TTATVLARSI) :::::: TTATVLAQALI	190 NIISDAMKKVC	: :: :: DLIAEAMDKVC	260	IQSIVPALEIA : :: VKDLLPLLEKV	330	VFGEEGLTLNL: :: :	400	LAKLSDGVAVL	CAKLAGGVAVI
40	ADARALMLQG :: EEARRGLERG	110	NTNEEAGDGT : :::: KTDDVAGDGT	180 , ISANGDKEIG	::::::	250	7LLSEKKISS ::::::::::::::::::::::::::::::::::	320	DMAIATGGA' :::::::	390	YEKEKLNERI	YDREKLQERI
30	AAYAKDVKFG. :: AKTIAYD	100	AKLVQDVANI : :: AELVKEVAKI	170 PEEIAQVATI	KEQIAATAAI	240	QKCEFQDAYV : QEAVLEDPYI	310	FGDNRKNOLK::::::::	380	IEQLDVTTSE	RQEIENSDSD
20	/SRVLAPHLTI	06	DLKDKYKNIG : : : : :	160 LKKQSKPVTT	:::::	230	SPYFINTSKG : :: SGYFVTDPER	300	LQVVAVKAPG :::::: FKSVAVKAPG	370	AQIEKRIQEI	DAIAGRVAQI
10	MLRLPTVFRQMRPVSRVLAPHLTRAYAKDVKFGADARALMLQGVDLLADAVAVTMGPKGRTVIIEQSWGS : :::::::::::::::::::::::::::::::::::	80	PKVTKDGVTVAKSIDLKDKYKNIGAKLVQDVANNTNEEAGDGTTTATVLARSIAKEGFEKISKGANPVEI : :::::::::::::::::::::::::::::::::::	1 , RRGVMLAVDAVIAELKK	:: :: :: :: :: : : : : : : : : : : : :	220	LEIIEGMKFDRGYISPYFINTSKGQKCEFQDAYVLLSEKKISSIQSIVPALEIANAHRKPLVIIAEDVDG :: ::: :: :: :: :: :: :: :: :: :: :: ::	290	EALSTLVLNRLKVGLQVVAVKAPGFGDNRKNQLKDMAIATGGAVFGEEGLTLNLEDVQPHDLGKVGEVIV :::::::::::::::::::::::::::::::::::	360	TKDDAMLLKGKGDKAQIEKRIQEIIEQLDVTTSEYEKEKLNERLAKLSDGVAVLKVGGTSDVEVNEKKDR	TKDETTÌVEGAGDTDAIAGRVAQIRQEIENSDSDYDREKLQERLAKLAGGVAVIKAGAATEVELKERKHR
-		71		14		211		281		351		
	HUMP1 TB65K		HUMP1 TB65K	HUMP1	TB65K		HUMP1 TB65K		HUMP1 TB65K		HUMP1	TB65K

na The Shares

13. 18. 1.

S. C. S. S. A. Mar.

SECTAINAGES.

A STANDARD STAND

## FIGURE 4 CONT'D

490	GVEGSLI	GLEPGVV	560	KDPGMGA	KASVPG-					3.16
480	AMTIAKNA	LKQIAFNS	550	VTEIPKEE	VADKPEKE					Mean = 3413.16
470	SIEIIKRTLKIP	SANIVKVALEAP	540	VASLLTTAEVV	: : : : : : : : : : : : : : : : : : :				residues	23.23 Me
460	PANEDOKIC:	K-LEGDEATC	530	TRTALLDAAG	DPVKVTRSALQNAASI	•			s between	ation =
450	VTDALNATRAAVEEGIVLGGGCALLRCIPALDSLTPANEDQKIGIEIIKRTLKIPAMTIAKNAGVEGSLI	TLLQAA	520	VEKIMQSSSEVGYDAMAGDFVNMVEKGIIDPTKVVRTALLDAAGVASLLTTAEVVVTEIPKEEKDPGMGA	AEKVRNLPAGHGLNAQTGVYEDLLAAGVADPVKVTRSALQNAASIAGLFLTTEAVVADKPEKEKASVPG-				eaks 540 possible matches between residues	Standard deviation
440	EGIVLGGGC	EGIVAGGGV	510	DAMAGDEVN	NAQTGVYED				breaks f 540 pos	16 SD
430	LNATRAAVEE	VRNAKAAVE	500	MQSSSEVGY	RNLPAGHGL	570	MGGMGGGMGGGMF	GGDMGGMDF	score = 4560, 5 breaks identities out of 540	s re = 49.3
421	VTDA::	IEDA	491	VEKI	AEKV	561	MGGM		score = identit	25 random runs Alignment score =
	HUMP1	TB65K		HUMP1	TB65K		HUMP1	TB65K	Total 257	25 ran

THE STATE OF THE S 

## INTERNATIONAL SEARCH REPORT

International Application No PCT/US 89/02619

I. CLAS	SIFICATI	N OF SUBJECT MATTER (It several cit	Assistantian symbols apply indicate all \$	703 03/02019
Accordin	ng to Internal	tional Patent Classification (IPC) or to both	National Classification and IPC	
IPC <sup>4</sup> :		K 37/02, A 61 K 39/0		
II. FIELE	S SEARCE	HED		
		Minimum Docu	mentation Searched 7	
Classifica	tion System		Classification Symbols	
IPC4	· · ·	C 12 N, A 61 K		
		Documentation Searched oth to the Extent that such Docume	er than Minimum Documentation ints are included in the Fields Searched	
III. DOC	UMENTS C	ONSIDERED TO BE RELEVANT		
Category *	Citat	ion of Document, 15 with Indication, where a	appropriate, of the relevant passages 12	Relevant to Claim No. 13
X	EP,	A, 0262170 (DE STAAT 6 April 1988, see the whole docume	DER NEDERLANDEN) -	1-5
x	]	A, 88/00974 (WHITEHE BIOMEDICAL RESEARCH) see claims 8-17	AD INSTITUTE FOR 11 February 1988,	1,2,6,7
	I	Natl. Acad. Sci, US June 1988, (US) D. Young et al.: "Str immune targets in ler tuberculosis", pages see the whole article	ress proteins are prosy and 4267-4270,	1-7
Х	I I I I I I I I I I I I I I I I I I I	Natl. Acad. Sci, USA March 1987 (US) R.N. Husson et al.: " major protein antiger suberculosis: The eti suberculosis and lepr mmunodominant antige	Genes for the solutions of Mycobacterium cologic agents of solutions on the solutions of th	1,2,6,7
"A" doct contain the contain t	ument definire sidered to be ser document grate ument which is cited to ion or other ument referring means ument publish than the profile of than the profile in the profil	of cited documents: 19 Ing the general state of the art which is not of particular relevance but published on or after the international may throw doubts on priority claim(s) or establish the publication date of another apecial reason (as specified) ing to an oral disclosure, use, exhibition or ned prior to the international filing date but outly date claimed pletion of the international Search	"T" later document published after the or priority date and not in conflictive to understand the principle invention.  "X" document of particular relevance cannot be considered novel or clinvolve an inventive step.  "Y" document of particular relevance cannot be considered to involve a document is combined with one of ments, such combination being of in the art.  "4" document member of the same particular in the art.	t with the application but or theory underlying the e; the claimed invention cannot be considered to e; the claimed invention inventive step when the ir more other such docu-vious to a person skilled tent family
			Date of Mailing of this International Seat 2 ft. 10, 89	rch Report
	Septe	mber 1989		
		Authority	Signature of Authorized Officer	

Calegory .	Citation of Doguests	1,2,6,7  1,2,6,7  1,2,6,7  1,2,6,7
	Citation of Document, with indication, where appropriate, of the relevant passages	
j		i Nevent to Claim No
!	D2G07 1670 1602	
į	Pages 1679-1683,	
İ	see the whole article	
. X	Nature, vol. 316, 1 August 1985,	
i	R.A. Young et al.: "Genes for the	1,2,6,7
:	major protein antigens of the	
· .:	major protein antigens of the leprosy parasite Mycobacterium leprae", pages 450-452.	
1		·
į	see the whole article	
$\mathbf{x}^{\pm}$	Infoction con	İ
* :	Infection & Immunity, vol. 55, no. 6, 1987, American Soc. for Microbiology (MS)	1 2 6 7
:	American Soc. for Microbiology (US)	-,2,0,1
	A.M.W. THOTE Et 31 "Characteria"	
	genicity of a 64-kilodalter muno-	
İ	of Mycobacterium bovis BCG expressed	
!	docher tenta coli k=17"	
İ	Pages 1466-1475	
	see the whole article	
A	WO 3 05/05004	
	WO, A, 85/05034 (UNIVERSITY COLLEGE,	1-5
	LONDON) 21 November 1985, see claims	- 0
P,X	WO, A, 88/05823 (WHITEHEAD INSTITUTE FOR	
	MEDICAL RESEARCH) 11 August 1988,	1,2,6,7
. !	see the whole document	
?,X;		;
. ,	WO, A, 88/06591 (SCRIPPS CLINIC AND	1 2 6 7
	THOUGHT FUINITITION I	1,2,0,1
	7 September 1988, see claims	5
		:
İ		
•		. [
	a single receipt the first and and	
:	ig bions un romalismitui	
	round with a pennishing transport takes "The Control of temperature and the Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of Control of C	onida Aspra I
and the same	and the state of the confidence of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state of the state	Televist 18th
. •		THE ME NEW TOTAL
	and the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of the same of th	CAN PRINCE
:		
i		
:		
	1	

PORTAR	INFORMATI	ON CONTIN	UED FROM TH	E SECOND S	HEET			
. 1							1	
}							<b> </b>	
ļ.							}	
ţ							ı	
							1	
[	••••							
		•	•	•				•
1		•		تواليموا والمهوا والو	ing the second of the second of the second of the second of the second of the second of the second of the second of the second of the second of the second of the second of the second of the second of the second of the second of the second of the second of the second of the second of the second of the second of the second of the second of the second of the second of the second of the second of the second of the second of the second of the second of the second of the second of the second of the second of the second of the second of the second of the second of the second of the second of the second of the second of the second of the second of the second of the second of the second of the second of the second of the second of the second of the second of the second of the second of the second of the second of the second of the second of the second of the second of the second of the second of the second of the second of the second of the second of the second of the second of the second of the second of the second of the second of the second of the second of the second of the second of the second of the second of the second of the second of the second of the second of the second of the second of the second of the second of the second of the second of the second of the second of the second of the second of the second of the second of the second of the second of the second of the second of the second of the second of the second of the second of the second of the second of the second of the second of the second of the second of the second of the second of the second of the second of the second of the second of the second of the second of the second of the second of the second of the second of the second of the second of the second of the second of the second of the second of the second of the second of the second of the second of the second of the second of the second of the second of the second of the second of the second of the second of the second of the second of the second of the second of the second of the second o	والمعتدد والميتات		
·		and the second of the second			a an american	:	· .	·
·   :			•		•	•	1	• •
	•• •							
· ]			• •				İ	
			•.	-		and the second second second		
						<u> </u>	ŀ	
V. W OBSE	RVATIONS	WHERE CERT	TAIN CLAIMS V	WERE FOUND	UNSEARCH	ABLE		
This Internati	onal search re	port has not be	en established in	respect of cost	in alalma	Article 17(2) (a) fo		
1. Claim n	umbers 8 - 1	T because they	telete to subject	managed of Cert	in caims unger	ched by this Autho	r the following	reasons:
			relate to soplect	matter not req	uired to be sean	ched by this Autho	rity, namely:	
See	מי פוז	le 39.1	/ T171		•			
Moth	ror Ru	16 33.1	(IV)					•
CUTO	ious io.	r treat	ment of	the hun	nan or a	nimal boo	ly by	
Surg	LETA OT	therap	Y, as we	ll as c	liagnost	ic method	is.	••
	*			7.	<b>.</b> .		7 m 1 m	
•	ī.					والمساوم فتشكون والمتعاور والاستا		
Claim nu		. because they	relate to parts of	the internation:	d application the	at do not comply w		٠.
ments to	such an exter	nt that no mean	ingful internation	al search can b	e carried out, so	it oo not comply w	ith the prescri	bed require-
							2.1	416
	•		• •					
		• • • •				•		
			-			*		
						• •		:
Claim nu	mhere i			-				:
PCT Rule	6.4/a).	secures may an	Dependent claim	is and are not o	irafted in accordi	ance with the secor	nd and third se	ntences of
			·		÷		•	
I. OBSER	VATIONS W	HERE UNITY	OF INVENTIO	N IS LACKU	16.2	<del></del>		
No Internation	-1.6			· · · · · · · · · · · · · · · · · · ·		<u></u>	·	
iis internation	ial Searching	Authority found	muitiple invention	ns in this interr	ational applicati	on as follows:		
				•				
				**				
		•		•				
		•						
-	٠							
As all requ	uired additional	i search fees we	ere timely paid by	the applicant, t	nis International	search report cove		
As only so	me of the regi	uired additional	search fees were	timely paid by	the applicant to	his international se		
TUOSA CIGILI	ns of the intern	national applicat	tion for which fee	s were paid, ap	ecifically claims	:	arch report c	overs only
٦.,								
No required	d additional se	arch fees were	timely paid by the	applicant. Cor	sequently, this i	international search		
mia inasulic	in iirst mentior	ned in the claim	18; It is covered b	y claim number	E:		raport is fer	stricted to
1	•							
invite paym	hable claims c	ould be searche	t noffe twodtliw bi	ustifying an ad	ditional fee, the	International Searce	thing Aidha	مرامية
ark on Protei	at						······A WALVOLI	ry ard not
		-						
= agaiti0:	nai search leer	A Wate Accomos	aniad bu analiass.					
No		· · · · · · · · · · · · · · · · · · ·	anied by applicant	's protest				}
No protest	accompanied t	he payment of a	anted by applicant additional search	l's protest. fees.				

## ANNEX TO THE INTERNATIONAL SEARCH REPORT ON INTERNATIONAL PATENT APPLICATION NO.

US 8902619

SA 29466

This annex lists the patent family members relating to the patent documents cited in the above-mentioned international search report. The members are as contained in the European Patent Office EDP file on 18/10/89 The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

Patent document cited in search report	Publication date		ent family ember(s)	Publication date
EP-A- 0262170	06-04-88	FR-A- WO-A- JP-T- US-A-	2596593 8706073 63502871 4821004	02-10-87 08-10-87 20-10-88 11-04-89
WO-A- 8800974	11-02-88	AU-A- EP-A-	7801387 0316354	24-02-88 24-05-89
WO-A- 8505034	21-11-85	AU-A- EP-A- JP-T- US-A-	4297685 0181364 61502258 4716038	28-11-85 21-05-86 09-10-86 29-12-87
WO-A- 8805823	11-08-88	AU-A-	1548388	24-08-88
WO-A- 8806591	07-09-88	AU-A- EP-A- ZA-A-	1495688 0305488 8801344	26-09-88 08-03-89 26-08-88

For more details about this annex : see Official Journal of the European Patent Office, No. 12/82